REFERENCES

1. Didier G, Stephane B, John S. Amoxicillin-Clavulanate Therapy Increases Childhood Nasal Colonization by Methicillin-Susceptible *Staphylococcus aureus* Strains Producing High Levels of Penicillinase. Antimicrobial. Agents Chemother, 2004; 48: 4618-4623.

2. Foster TJ, Hook M. Surface protein adhesins of Staphylococcus aureus. Trends Microbiol 1998; 6:484–8.

3. Jevons, M. Patricia. "Celbenin"- resistant Staphylococci. Br Med J. 1961 Jan 14; 15219:124-125. (PMC free article).

4. Hussain, F. M., Boyle-Vavra, S., Bethel, C. D. & Daum, R. S. *Pediatr. Infect. Dis. J.*2000; 19, 1163–1166.

5. Centers for Disease Control and Prevention *Morbid. Mortal. Wkly. Rep.* 1999; 48, 707–710.

6. Public Health Division: Provincial Methicillin-Resistant Staphylococcus aureus (MRSA) infection prevention and control guidelines Edmonton; 2007.

7. Chambers HF: Bacterial & chlamydial infections. In Current medical diagnosis and treatment... 47 editions. Edited by: McPhee SJ, Papadakis MA, Tierney LM. New York: McGraw-Hill; 2008:1232-1235.

8. Nicolle L: Community-acquired MRSA: a practitioner's guide. CMAJ 2006; 175:145-146.

9. Jarvis WR, Schlosser J, Chinn RY, Tweeten S, Jackson M. National prevalence of methicillin-resistant *Staphylococcus aureus* in inpatients at US health care facilities, Am. J. Infect. Control, 2007; 35: 631-637.

10. Kim T, Oh PI, Simor AE, The economic impact of methicillin resistant *Staphylococcus aureus* in Canadian hospitals. Infect.Control Hosp. Epidemiol., 2001; 22: 99-104. 13.

11. Lodise TP, McKinnon PS, Clinical and economic impact of methicillin resistance in patients with *Staphylococcus aureus* bacteremia. Diagn. Microbiol. Infect. Dis., 2005; 52: 113-122. 14.

12. Kearns AM, Seiders J, Wheeler R, Steward M, Rapid detection of methicillin resistant staphylococci by multiplex pCK .J. Hosp.Infect.1999; 43: 33-37.

13. National Nosocomial Infections Surveillanc, Systems Report, data summary from January 1992 through June 2004, issued October 2004 Am. J. Infect. Control, 2004; 32: 470-485.

14. Harbarth S, Martin Y, Rohner P, Henry N, Auckenthaler R, Pittet D, Effect of delayed infection control measures on a hospital outbreak of methicillin- resistant *Staphylococcus aureus*. J. Hosp. Infect., 2000; 46: 43-49.

15. Calfee DP, Durbin LJ, Germanson TP, Spread of methicillinresistant *Staphylococcus aureus* (MRSA) among household contacts of individuals with nosocomially acquired MRSA. Infect. Control Hosp. Epidemiol., 2003; 24: 422 -426.

16. Wannet W, Heck M, Pluister G, Spalburg E, De Neeling AJ, Panton-valentine leucocidine positive MRSA in 2003: the Duch situation. Eur. Surveillance2004; 9: 28-29.

17. Ibelings MMS, Bruining HA, Methicillin-resistant *Staphylococcus aureus*: acquisition and risk of death in patients in the intensive care unit. Eur. J. Surg., 1998; 164: 411 –1.

18. Law MR, Gill ON, Hospital acquired infection with methicillinresistant and methicillin sensitive staphylococci. Epidemiol.Infect, 1998; 101: 623 –9.

19. Pujol M, Pena C, Pallares R , Risk factors for nosocomial bacteraemia due to methicillin-resistant *Staphylococcus aureus.* Eur. J. Clin Microbiol., Infect. Dis, 1994; 13: 96 –100.

20. Merrer J, Santoli F, Vecchi C, Colonization pressure and risk of acquisition of methicillin resistant *Staphylococcus aureus* in a medical intensive care unit. Infect. Control. Hosp. Epidemiol, 2000; 21: 718 –23.

21. Girou E, Pujade G, Legrand P, Selective screening of carriers for control of methicillin-resistant *Staphylococcus aureus* (MRSA) in high-risk hospital areas with a high level of endemic MRSA. Clin. Infect Dis, 1998; 27: 543 –50.

22. Lucet JC, Chevret S, Durand-Zaluski I , Prevalence and risk factors for carriage of methicillin resistant *Staphylococcus aureus* at admission to the intensive care unit. Arch. Int. Med, 2003; 163: 181 –8.

23. Arnold MS, Dempsey JM, Fishman M, The best hospital practices for controlling methicillin resistant *Staphylococcus aureus* on the cutting edge. Infect. Control Hosp. Epidemiol, 2002; 23: 69 –76.

24. Hill RA, Duckworth GJ, Casewell M, Elimination of nasal carriage of methicillin resistant *Staphylococcus aureus* with mupirocin during a hospital outbreak. J. Antimicrob. Chemother, 1988; 22: 377 –84.

25. Rampling A, Wiseman S, Davis L, Evidence that hospital hygiene is important in the control of methicillin resistant *Staphylococcus aureus*. J. Hosp. Infect.,2001; 49: 109 – 116.

26. Deurenberg R H, Stobberingh E. E., The evolution of *Staphylococcus aureus* Infection, Genetics and Evolution. 2008; 751-753.

27. Hiramatsu, K., Hanaki, H., Ino, T., Yabuta, K., Oguri, T. & Tenover, F. C., *J. Antimicrob. Chemother, 1997;* 40, 135–136.

28. Centers for Disease Control and Prevention, *Morbid. Mortal. Wkly. Rep.* 1997; 46:765–766.

29. Barber M, Methicillin Resistant Staphylococci. *J. Clin. Pathol, 1961* 14:385-393. Center of Disease Control and Prevention. Four pediatric deaths from community-acquired methicillin resistant Staphylococcus aureus. JAMA 1999; 282:1123-1125.

30. Verma S, Joshi S, Chitnis V, Hemwani N, Chitnis D**,** Growing problems of Methicillin Resistance Staphylococci - Indian Scenario. *Indian J. Med Sci., 2000;* 54:535-540.

31. Prere MF, O Baron, S Cohen Bacrie, O Fayet, Genotype MRSA, a new genetic test for rapid identification of Staphylococci and detection of *mecA* gene. *Pathologie Biologie., 2006;* 54:502-505.

32. Moreillon P, Que Y A, Glauser M P .,Staphylococcus aureus(Including Staphylococcal Toxic Shock) Chapter 192 In Mandell G L., Bennett J E., Dolin RPriniples and Practice of Infecious Diseases vol- 2 6th ed. Elesevier Churchill Livingstone,2005 p 233.

33. Ajantha GS, Kulkarni RD, Shetty J, Shubhada C, Jain P. Phenotypic detection of inducible clindamycin resistance among *Staphylococcus aureus* isolates by using the lower limit of recommended inter-disk distance. *Indian J Pathol Microbiol*. 2008; 51:376–78.

34. Leclercq R. Mechanisms of resistance to macrolides and lincosamides: nature of the resistance elements and their clinical implications. *Clin Infect Dis*. 2002; 34:482–92.

35. Ross JI, Eady EA, Cove JH, Cunliffe WJ, Baumberg S, Wooton JC. Inducible erythromycin resistance in *staphylococci* is encoded by a member of the ATP-binding transport super-gene family. *Mol Microbiol*. 1990; 4:1207–14.

36. Ross JI, Farrell AM, Eady EA, Cove JH, Cunliffe WJ. Characterisation and molecular cloning of the novel macrolide–streptogramin B resistance det*erm*inant from Staphylococcus epid*erm*idis. *J Antimicrob Chemother*. 1989; 24:851–62.

37. Roberts MC, Sutcliffe J, Courvalin P, Jensen LB, Rood J, Seppala H. Nomenclature for macrolide-lincosamide–streptogramin B resistance det*erm*inants. *Antimicrob Agents Chemother*. 1999; 43:2823–30.

38. Lewis JS, Jorgensen JH. Inducible clindamycin resistance in staphylococci: should clinicians and microbiologists be concerned? *Clin Infect Dis*. 2005; 40:280–85.

39. Vivek JS, Rajesh GN, Mukesh S, Manpreet K, Manpreet K, Misra RN, et al. Prevalence of inducible clindamycin resistance among community- and hospital-associated *Staphylococcus aureus* isolates in a tertiary care hospital in India. Biomed Res. 2011; 22:465–9.

40. Saderi H, Emadi B, Owlia P. Phenotypic and genotypic study of macrolide, lincosamide and streptogramin B (MLSB) resistance in clinical isolates of *Staphylococcus aureus* in Tehran, Iran. Med Sci Monit. 2011; 17:BR48–53. [PMC free article] [PubMed]

41. Reddy SP, Suresh R. Phenotypic detection of inducible clindamycin resistance among the clinical isolates of *staphylococcus aureus* by using the lower limit of inter disk space. J Microbiol Biotechnol Res. 2012; 2:258–64.

42. Zelazny AM, Ferraro MJ, Glennen A, Hindler JF, Mann LM, Munro S, et al. Selection of strains for quality assessment of the disk induction method for detection of inducible clindamycin resistance in staphylococci: A CLSI collaborative study. J Clin Microbiol. 2005; 43:2613–5. [PMC free article][PubMed]

43. Seifi N, Kahani N, Askari E, Mahdipour S, Naderi NM. Inducible clindamycin resistance in *Staphylococcus aureus* isolates recovered from Mashhad, Iran. Iran J Microbiol. 2012;4:82–6.[PMC free article] [PubMed]

44. Steward CD, Raney PM, Morrell AK, Williams PP, McDougal LK, Jevitt L, et al. Testing for induction of clindamycin resistance in erythromycin-resistant isolates of *Staphylococcus aureus*. J Clin Microbiol. 2005; 43:1716–21. [PMC free article] [PubMed]

45. Aktas Z, Aridogan A, Kayacan CB, Aydin D. Resistance to macrolide, lincosamide and streptogramin antibiotics in staphylococci isolated in Istanbul, Turkey. J Microbiol. 2007; 45:286–90. [PubMed]

46. Schmitz FJ, Sadurski R, Kray A, Boos M, Geisel R, Köhrer K, et al. Prevalence of macrolide-resistance genes in *Staphylococcus aureus* and Enterococcus faecium isolates from 24 European university hospitals. J Antimicrob Chemother. 2000; 45:891–4. [PubMed]

47. Rajaduraipandi K, Mani KR, Panneerselvam K, Mani M, Bhaskar M, Manikandan P. Prevalence and antimicrobial susceptibility pattern of methicillin resistant *Staphylococcus aureus*: a multicentre study. *Indian J Med Microbiol*. 2006; 24: 34-38.

48. Shantala G B, Shetty AS, Rao RK et al. detection of inducible clindamycin resistance in clinical isolates of *Staphylococcus aureus* by the disc diffusion induction test. *J Clin Diag Res*. 2011; 5(1):35-37.

49. Smith TL, Pearson ML, Wilcox KL, Cruz C, Lancaster MV, Robinson-Dunn B, et al. Emergence of vancomycin resistance in *Staphylococcus aureus*. glycopeptide intermediate *Staphylococcus aureus* working group. *N Eng J Med*. 1999; 340: 493-501.
50. Naimi TS, LeDell KH, Como-Sabetti K, Borchardt SM, Boxrud DJ, Etienne J, et al. Comparison of community- and health-care associated methicillin resistant *Staphylococcus aureus* infection. *JAMA*. 2003; 290: 2976-84.

51. Lertcanawanichakul M, Chawawisit K, Choopan A, Nakbud K, Dawveera Kul K. Incidence of constitutive and inducible clindamycin resistance in clinical isolates of methicillin resistant *Staphylococcus aureus. Walailak J Sci & Tech.* 2007; 4:155-63.

52. Surendra kumar; *Staphylococcus* .Textbook of Microbiolgy 2012. Jaype Brothers Medical Publishers (P) Ltd.pp 229-239.

53. Sharon J. P, *Staphylococcus.* Topley & Wilson's Microbiology & Microbial Infections Vol- 2t 10th ed. Hodder Arnold p.778.

54. Winn W C, Allen S D, Janda W M, Koneman E W, Procop G W, Schreckenberger P C, et al Gram Positive Cocci Part 1: *Staphylococci* and related Gram Positive Cocci, Colr Atlas and Textbook of Diagnostic Microbiology 6th ed Lippincott Williams & Wilkins p. 645.

55. Forbes B A., Sahm D F, Weissfeld A S., et al, Part 3rd Bacteriology, Staphylococcus, Micrococcus, and Similar organisms Bailey & Scott's Diagnostic Microbiology 12 ed. Mosby Elsevier p. 258.

56. Ananthanarayan R, Paniker C K J. Staphylococcus. Textbook of Microbiology 9 ed. Universities Press (India) Private Limited.

57. Duran N, Ozer B, Duran G G , Onlen Y, Demir C. Antibiotic resistance genes & susceptibility patterns in staphylococci. Indian J of Med Res 2012; 135(3): 389-396.

58. Pal N, Sharma B, Sharma R, Vyas. L Detection of inducible Clindamycin resistance among Staphylococcal isolates from different clinical specimens in western India. J Postgrad Med. 2010; 56:182-5.

59. Laclercq R. Mechanisms of resistance to macrolides and lincosamides: Nature of resistance elements and their clinical implications. Clin Infect Dis 2002; 34:482-92.

60. Drinkovic D, Fuller ER, Shore KP, Holland DJ, Ellis-Pegler R. Clindamycin treatment of *Staphylococcus aureus* expressing inducible clindamycin resistance. J Antimicrob Chemother 2001; 48:315-6.

61. Fiebelkorn KR, Crawford SA, McElmeel ML, Jorgensen JH. Practical disk diffusion method for detection of inducible clindamycin resistance in *Staphylococcus aureus* and coagulase negative staphylococci. J Clin Microbiol 2003; 41:4740-4.

62. Mehndiratta PL, Bhalla P. typing of methicillin resistant Staphylococcus aureus. A technical review. Indian J Med Microbiol 2012; 30: 16-23.

63. Mathur SK, Singhal S, Prasad KN, Kishore J, Ayyagari A. Prevalence of methicillin resistant *Staphyloccous aureus* in a tertiary care hospital. Indian J Med Microbiol 1994; *12* : 96-101.

64. Pal N, Ayyagari A. Drug resistance pattern of methicillin resistant *Staphylococcus aureus.* Indian Paediatr 1991; *28* : 725-7.

65. Mohanasoundaram K M, Lalitha M.K. Comparison of phenotypic versus genotypic methods in the detection of methicillin resistance in *Staphylococcus aureus.* Indian J Med Res 2008; 127(1): 78-84.

66. Cai Yongwe, Kong F, Wang Q, Tong Z, Sintehenko V, ZengX, et al. Comparison of single and multilocus sequence typing and toxin gene profileing for characterization of methicillin resistant staphylococcus aureus. J Clin Microbiol 2007; 45:3302-8.

67. Cree RG, Aleljung P, Paulsson M. Cell surface hydrophobicity and adherence to extra cellular matrix proteins in two collections of methicillin resistant Staphylococcus aureus. J Med Microbiol 1992; 36: 164-71.

68. Vaudaux PE, Monzillo V, Francois P. Introduction to the mec element (Methicillin resistance) into Staphylococcus aureus alters in vitro functional activities of fibrogen and fibronectin adhesions. Antimicrob Agents Chemother 1998; 42:564-70.

69. Sharma NK, Rees CE, Dodd CE. Development of a single reaction, multiplex PCR toxin typing assay for staphylococcus strains. J Appl Environ Microbiol 2006; 66: 1347-53.

70. Cameron DR, Howden BP, Peleg AY. The interface between antibiotic Resistance and Virulence in Staphylococcus aureus and its Impact upon Clinical Outcomes. Clinical infectious Dis. 2011; 53 (6): 576-582.

71. Berger-Bachi B, Rohrer S. Factors influencing methicillin resistance in staphylococci. Arch Microbiol 2002; 178: 165–171.

72. Katayama Y, Ito T, Hiramatsu K. Genetic organization of the chromosome region surrounding mecA in clinical staphylococcal strains: role of IS431-mediated mecI deletion in expression of resistance in mecA-carrying, lowlevel methicillin-resistant *Staphylococcus haemolyticus*. Antimicrob Agents Chemother 2001; 45: 1955–1963.

73. Rosato AE, Kreiswirth BN, Craig WA, Eisner W, Climo MW, Archer GL. *mecA-blaZ* Corepressors in Clinical *Staphylococcus aureus* Isolates. Antimicrobial agents and chemotherapy, 2003; 47(4): 1460–1463.

74. Lowy FD. Antimicrobial resistance: the example of Staphylococcus aureus J. Clin. Invest, 2003; 111:1265–1273.

75. Deurenberg R. H., Vink C., Kalenic S., Friedrich A. W., Bruggeman C. A. and Stobberingh E. E. The molecular evolution of methicillin-resistant *Staphylococcus aureus*. Clin Microbiol Infect 2007; 13: 222–235.

76. Ito T, Katayama Y, Asada K et al. Structural comparison of three types of staphylococcal cassette chromosome mec integrated in the chromosome in methicillin-resistant *Staphylococcus aureus*. Antimicrob Agents Chemother 2001; 45: 1323–1336.

77. Daum RS, Ito T, Hiramatsu K et al. A novel methicillin resistance cassette in community-

acquired methicillin-resistant *Staphylococcus aureus* isolates of diverse genetic backgrounds. J Infect Dis 2002; 186: 1344–1347.

78. Wielders CLC, FluitAC, Brisse S, Verhoef J, Schmitz FJ. *mecA* Gene Is Widely Disseminated in *Staphylococcus aureus* Population. Journal of Clinical Microbiology, 2002; 40(11): 3970–3975.

79. Pramodhini S, Thenmozhivalli PR, Selvi R, Dillirani V, Vasumathi A, Agatha. Comparison of Various Phenotypic Methods and *mecA* Based PCR for the Detection of MRSA. Journal of Clinical and Diagnostic Research. 2011; 5(7): 1359-1362.

80. Leclercq R. Mechanisms of resistance to macrolides and lincosamides: nature of the resistance elements and their clinical implications. *Clin Infect Dis*. 2002; 34: 482-92.

81. Lewis JS 2nd, Jorgensen JH. Inducible clindamycin resistance in *Staphylococci*: should clinicians and microbiologists be concerned? *Clin Infect Dis*. 2005; 40: 280-85.

82. Levin TP, Suh B, Axelrod P, Truant AL, Fekete T. Potential clindamycin resistance in clindamycin-susceptible, erythromycin-resistant *Staphylococcus aureus*: report of a clinical failure. *Antimicrob Agents Chemother*. 2005; 49: 1222-24.

83. Rao GG. Should clindamycin be used in treatment of patients with infections caused by erythromycin-resistant *staphylococci*? *J Antimicrob Chemother*. 2000; 45: 715.

84. Drinkovic D, Fuller ER, Shore KP, Holland DJ, Ellis-Pegler R. Clindamycin treatment of *Staphylococcus aureus* expressing inducible clindamycin resistance. *J Antimicrob Chemother*. 2001; 48: 315-16.

85. Delialioglu N, Aslan G, Ozturk C, Baki V, Sen S, Emekdas G. Inducible clindamycin

resistance in *Staphylococci* isolated from clinical samples. *Jpn J Infect Dis*. 2005; 58:104-06.

86. Fiebelkorn KR, Crawford SA, McElmeel ML, Jorgensen JH. Practical disk diffusion method for detection of inducible clindamycin resistance in *Staphylococcus aureus* and coagulase-negative *staphylococci*. *J Clin Microbiol.* 2003; 41: 4740-44.

87. Yilmaz G, Aydin K, Iskender S, Caylan R, Koksal I. Detection and prevalence of inducible resistance in *Staphylococci*. *J Med Microbiol.* 2007; 56:342-45.

88. Vivek JS, Rajesh GN, Mukesh S, Manpreet K, Manpreet K, Misra RN, et al. Prevalence of inducible clindamycin resistance among community- and hospital-associated *Staphylococcus aureus* isolates in a tertiary care hospital in India. Biomed Res. 2011; 22:465–9.

89. Saderi H, Emadi B, Owlia P. Phenotypic and genotypic study of macrolide, lincosamide and streptogramin B (MLSB) resistance in clinical isolates of *Staphylococcus aureus* in Tehran, Iran. Med Sci Monit. 2011;17:BR48–53. [PMC free article] [PubMed]

90. Reddy SP, Suresh R. Phenotypic detection of inducible clindamycin resistance among the clinical isolates of *staphylococcus aureus* by using the lower limit of inter disk space. J Microbiol Biotechnol Res. 2012;2:258–64.

91. Zelazny AM, Ferraro MJ, Glennen A, Hindler JF, Mann LM, Munro S, et al. Selection of strains for quality assessment of the disk induction method for detection of inducible clindamycin resistance in staphylococci: A CLSI collaborative study. J Clin Microbiol. 2005;43:2613–5. [PMC free article][PubMed]

92. Seifi N, Kahani N, Askari E, Mahdipour S, Naderi NM. Inducible clindamycin resistance in *Staphylococcus aureus* isolates recovered from Mashhad, Iran. Iran J Microbiol. 2012;4:82–6.[PMC free article] [PubMed]

93. Steward CD, Raney PM, Morrell AK, Williams PP, McDougal LK, Jevitt L, et al. Testing for induction of clindamycin resistance in erythromycin-resistant isolates of *Staphylococcus aureus*. J Clin Microbiol. 2005; 43:1716–21. [PMC free article] [PubMed]

94. Aktas Z, Aridogan A, Kayacan CB, Aydin D. Resistance to macrolide, lincosamide and streptogramin antibiotics in staphylococci isolated in Istanbul, Turkey. J Microbiol. 2007; 45:286–

90. [PubMed]

95. Schmitz FJ, Sadurski R, Kray A, Boos M, Geisel R, Köhrer K, et al. Prevalence of macrolide-resistance genes in *Staphylococcus aureus* and Enterococcus faecium isolates from 24 European university hospitals. J Antimicrob Chemother. 2000;45:891–4. [PubMed]

96. Fiebelkorn KR, Crawford SA, McElmeel ML, Jorgensen JH. Practical disk diffusion method for detection of inducible clindamycin resistance in *Staphylococcus aureus* and coagulase-negative staphylococci. J Clin Microbiol. 2003; 41:4740–4. [PMC free article] [PubMed]

97. Coutinho Vde L, Paiva RM, Reiter KC, de-Paris F, Barth AL, Machado AB. Distribution of *erm* genes and low prevalence of inducible resistance to clindamycin among staphylococci isolates. Braz J Infect Dis. 2010;14:564–8. [PubMed]

98. Sedighi I, Yousefi Mashouf R, Pak N, Seif Rabiee MA. D-test method for detection of inducible clindamycin resistance in *Staphylococcus aureus*. Iran J Pediatr. 2009;19:293–7.

99. O'Sullivan MV, Cai Y, Kong F, Zeng X, Gilbert GL. Influence of disk separation distance on accuracy of the disk approximation test for detection of inducible clindamycin resistance in *Staphylococcus spp*. J Clin Microbiol. 2006;44:4072–6. [PMC free article] [PubMed]

100. Jevons MP. 'Celbenin'-resistant Staphylococci. BMJ. 1961;1:124.

101. Patel AK, Patel KK, Patel KR, Shah S, Dileep P. Time trends in the epidemiology of microbial infections at a tertiary care center in west India over last 5 years. J Assoc Physicians India. 2010;58(Suppl):37–40.[PubMed]

102. Gopalakrishnan R, Sureshkumar D. Changing trends in antimicrobial susceptibility and hospital acquired infections over an 8 year period in a tertiary care hospital in relation to introduction of an infection control programme. J Assoc Physicians India. 2010;58(Suppl):25–31. [PubMed]

103. D'Souza N, Rodrigues C, Mehta A. Molecular characterization of Methicillin-resistant *Staphylococcus aureus* with emergence of epidemic clones of sequence type (ST)22 and ST 772 in Mumbai, India. J Clin Microbiol. 2010;48:1806–11. [PMC free

article] [PubMed]

104. Anupurba S, Sen MR, Nath G, Sharma BM, Gulati AK, Mohapatra TM. Prevalence of methicillin resistant Staphylococcus aureus in a tertiary referral hospital in eastern Uttar Pradesh. Indian J Med Microbiology. 2003; 21: 49-51.

105. Salaria M, Singh M. Methicillin resistant Staphylococcus aureus. Indian Pediatrics. 2001; 38:

29-36.

106. Pai V, Rao IR, Rao SP. Prevalence and Antimicrobial Susceptibility Pattern of Methicillin-resistant Staphylococcus Aureus [MRSA] Isolates at a Tertiary Care Hospital in Mangalore, South India. Journal of Laboratory Physicians. 2010; 2(2): 82-84.

107. David MZ and Daum RS. Community-Associated Methicillin-Resistant Staphylococcus aureus: Epidemiology and Clinical Consequences of an Emerging Epidemic. Clinical microbiology review. 2010; 23(3): 616–687.

108. Tiemersma EW, Bronzwear LAM, Lyytikainen O, degener JE, Schrijnemakers P, Bruinsma N, Monan J, Witte W, Grundmaun H. Methicillin resistant Staphylococcus aureus in Europe 1999-2002. • www.cdc.gov/eid • Emerge infect Disease. 2004; 10(9):1627-1634.

109. Borg M, Scicluna E, De Kraker M, Van de Sande-Bru insma N, Tiemersma E, Gur D et al. Antibiotic resistance in the southeastern Mediterranean - preliminary results from the AR Medical project (Staphylococcus aureus, European Surveillance). 2006; 20(11): 7.

110. Salgado CD, Farr BM, Calfee DP. Community acquired methicillin resistant Staphylococcus aureus: A meta-analysis of prevalence and risk factors. Clin. Infectious Disease. 2003; 131-139.

111. Verma S, Joshi S, Chitnis V, Hemwani N, Chitnis D. Growing problem of methicillin resistant staphylococci - Indian scenario. Indian J Med Sciences. 2000; 54:535-40.

112. Mathur SK, Singal, S, Prasad, KN, Kishore J, Ayyagiri A. Prevalence of MRSA in tertiary care hospital. Indian J Medical Microbiology. 1994;12: 96-101.

113. Tiwari HR, Sapkota D, Sen MR. High prevalence of multidrug-resistant MRSA in a tertiary care hospital of northern India. Infection and Drug Resistance. 2008; 1: 57–61.

114. Ganguly NK, Arora NK, Chandy SJ, Fairoze MN, Gill JP, Gupta U, *et al*. Rationalizing antibiotic use to limit antibiotic resistance in India. Indian J Med Res 2011; 134:281-94. [PUBMED] [Full text]

115. Indian Network for Surveillance of Antimicrobial Resistance (INSAR) group, India. Methicillin resistant *Staphylococcus aureus* (MRSA) in India: Prevalence & susceptibility pattern. Indian J Med Res 2013; 137:363-9.

116. Sader HS, Farrell DJ, Jones RN. Antimicrobial susceptibility of Gram-positive cocci isolated from skin and skin-structure infections in European medical centres. Int J Antimicrob Agents

2010; 36:28-32.

117. Daurel C, Huet C, Dhalluin A, Bes M, Etienne J, Leclercq R. Differences in potential for selection of clindamycin-resistant mutants between inducible erm(A) and erm(C) *Staphylococcus aureus* genes. J Clin Microbiol, 2008; 46: 546-550.

118. Herbert S, Barry P, Novick R.P. Subinhibitory clindamycin differentially inhibits transcription of exoprotein genes in *Staphylococcus aureus*. Infect Immun, 2001; 69: 2996-3003.

119. Marcinak J.F, Frank A.L. Epidemiology and treatment of community-associated methicillin-resistant *Staphylococcus aureus* in children. Expert Rev Anti Infect Ther,2006; 4: 91-100.

120. Jenssen W.D, Thakker-Varia S, Dubin D.T, Weinstein M.P.Prevalence of macrolides-lincosamides-streptogramin B resistance and erm gene classes among clinical strains of *staphylococci* and *streptococci*. Antimicrob Agents Chemother, 1987; 31: 883-888.

121. Braun L, Craft D, Williams R, Tuamokumo F, Ottolini M. Increasing clindamycin resistance among methicillin-resistant *Staphylococcus aureus* in 57 northeast United States military treatment facilities. Pediatr Infect Dis J, 2005; 24: 622-626.

122. Delialioglu N, Aslan G, Ozturk C, Baki V, Sen S, Emekdas G. Inducible clindamycin resistance in *staphylococci* isolated from clinical samples. Jpn J Infect Dis, 2005; 58: 104-106.

123. Fokas S, Tsironi M, Kalkani M, Dionysopouloy M. Prevalence of inducible clindamycin resistance in macrolide-resistant *Staphylococcus* spp. Clin Microbiol Infect ,2005; 11: 337-340.

124. Goyal R, Singh N.P, Manchanda V, Mathur M. (2004) Detection of clindamycin susceptibility in macrolide resistant phenotypes of *Staphylococcus aureus*. Indian J Med Microbiol2004; 22: 251-254.

125. Gadepalli R, Dhawan B, Mohanty S, Kapil A, Das B.K, Chaudhry R.Inducible clindamycin resistance in clinical isolates of *Staphylococcus aureus*. Indian J Med Res, 2006; 123: 571-573.

126. Navaneeth B.V. A preliminary in vitro study on inducible and constitutive clindamycin resistance in *Staphylococcus aureus* from a South Indian tertiary care hospital. Int J Infect Dis, 2006; 10: 184- 185.

127. Angel M.R, Balaji V, Prakash J, Brahmadathan K.N, Mathews M.S. Prevalence of inducible clindamycin resistance in gram positive organisms in a tertiary care centre. Indian J Med Microbiol, 2008; 26: 262-264.

128. Ciraj A.M, Vinod P, Sreejith G, Rajani K. Inducible clindamycin resistance among clinical isolates of *Staphylococci*. Indian J Pathol Microbiol, 2009; 52: 49-51.

129. Schmitz F.J, Petridou J, Fluit A.C, Hadding U, Peters G, von Eiff C, Distribution of macrolide-resistance genes in *Staphylococcus aureus* blood-culture isolates from fifteen German university hospitals. M.A.R.S. Study Group. Multicentre Study on Antibiotic Resistance in *Staphylococci*. Eur J Clin Microbiol Infect Dis,2000; 19: 385-387.

130. Gopal Rao G.Should clindamycin be used in treatment of patients with infections caused by erythromycin-resistant *staphylococci*? Journal of Antimicrobial Chemotherapy, 2000; 45: 715.

131. Drinkovic D, Fuller E.R, Shore K.P, Holland D.J, Ellis-Pegler R. Clindamycin treatment of *Staphylococcus aureus* expressing inducible clindamycin resistance. J Antimicrob Chemother, 2001; 48: 315-316.

132. Frank A.L et al. Clindamycin treatment of methicillin-resistant *Staphylococcus aureus* infections in children. Pediatr Infect Dis J, 2002; 21: 530-534.

133. Fiebelkorn K.R, Crawford S.A, McElmeel M.L, Jorgensen J.H.Practical disk diffusion method for detection of inducible clindamycin resistance in *Staphylococcus aureus* and coagulase-negative *staphylococci*. J Clin Microbio, 2003;l 41: 4740-4744.

134. Siberry G.K, Tekle T, Carroll K, Dick J. Failure of clindamycin treatment of methicillin-resistant *Staphylococcus aureus* expressing inducible clindamycin resistance in vitro. Clin Infect Dis, 2003; 37: 1257-1260.

135. Weisblum B, Siddhikol C, Lai CJ, Demohn V. Erythromycin-inducible resistance in *Staphylococcus aureus*: requirements for induction. J Bacteriol, 1971; 106: 835-847.

136. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial Susceptibility testing, Seventeenth informational supplement. Wayne PA.USA:CLSI2007.PP.1-182.CLSIdocumentM100-517.

137. Mathur P. Hand hygiene: Back to the basics of infection control. Indian J Med Res 2011; 134: 611-20.

138. Mathai E, Allegranzi B, Kilpatric C, Pittet D. prevention and control of health care – associated infections through improved hand hygiene. Indian J of Medical Microbiology 2011; 28(2):100-6.

139. WHO Guidelines on hand Hygiene in Health care. (World Health organization, Geneva) 2009. Available from: http://whqlibdoc.who.int/publications/2009.

140. Collee G J., Marmion B P., Fraser A G., Simmons A.,Staphylococcus: Cluster- forming Gram positive cocci,Chapter 11 Bacteria And Related Organisms, Mackie & McCartney Practical Medical Microbiology14th ed, Elesevier Churchill Livingstone.

141. Clinical and Laboratory Standard Institute, Performance Standards for Antimicrobial

Susceptibility Testing; Twenty Seven informational supplement. Wayne PA.USA:CLSI2017.PP.1-282.CLSIdocumentM 100- S27.

142. Sureerat C, Varaporn L, Manthana P, Urairat K, Sineenart K. Detection of inducible Clindamycin Resistance in Staphylococci by Disc Diffusion induction test. J Med Assoc Thai; 2009; 92(7): 947-51.

143. Qureshi A, Ingle R, MusaddiqM, Ali Y, Khan. Status and distribution of *mecA* gene in hospitalized patient's mrsa isolates. Bioscience Discovery, 2012; 3(1):52-57.

144. Pillai MM, Latha R, Sarkar G. Detection of Methicillin Resistance in *Staphylococcus aureus* by polymerase chain reaction and conventional methods: A Comparative Study. J Lab Physicians 2012; 4:83-8.